



Application of satellite surface wind data to ocean wind analysis and numerical weather prediction



R. Atlas, J. Ardizzone, R.N Hoffman,
M. Leidner, S. Majumdar, and J. C.
Jusem

Goals



- Produce consistent oceanic surface wind data of high quality and high temporal and spatial resolution for atmospheric and oceanic research and for improved weather and short-term climate prediction. – Generation of the Cross-Calibrated MultiPlatform (CCMP) Ocean surface wind data set.
- Evaluate (and enhance) the impact of each type of satellite surface winds on ocean surface wind analyses and nwp.



PO.DAAC

Physical Oceanography DAAC

Managing Data to Enable Understanding and Stewardship of the Ocean

ABOUT PO.DAAC

HOME

DATA CATALOG

TOOLS & SERVICES

RELATED SITES

ANNOUNCEMENTS

HELP/FAQ

Cross-Calibrated Multi-Platform (CCMP) Ocean Surface Wind Components

Attention Users: CCMP is now available on Public FTP.

MEaSURES - Making Earth Science Data Records for Use in Research Environments




[Products](#) / [Announcements](#) / [Citations/References](#) / [Related Publications](#)

Product Description

In collaboration with private and government institutions, a team led by Dr. Robert Atlas (PI; proposal originally solicited by REASoN, and currently funded by MEaSURES through NASA) has created a cross-calibrated, multi-platform (CCMP), multi-instrument ocean surface wind velocity data set (<http://sivo.gsfc.nasa.gov/oceanwinds>), for the period extending from January 1, 1987 through June 30, 2008, with wide ranging research applications in meteorology and oceanography. This product was a result of an investigation funded by NASA's Making Earth Science data records for Use in Research Environments (MEaSURES) Program.

It represents a continuation and expansion of the SSM/I surface




Quikscat and TRMM measurements over Hurricane Floyd on September 13, 1999.

Inputs to CCMP data set



- Cross-calibrated data sets from Remote Sensing Systems
 - SSM/I, TRMM TMI, AMSRE, Quikscat, Seawinds
- In situ data
 - Conventional ships and buoys
 - Tropical Atmosphere Ocean Project (TAO) buoys
 - Pilot Research Moored Array in the Atlantic (PIRATA) buoys
- Background analyses consist of the ECMWF operational analysis (from 1999) and the ERA-40 reanalysis
 - ECMWF operational analyses are available within 6-months of real-time at 1-degree resolution.
 - ERA-40 reanalyses are available through June 2002 at a resolution of approximately 2-degrees and are more accurate than ops. thru 1998.

Methodology




The variational analysis method (VAM) generates a gridded surface wind analysis which minimizes an objective function (J) measuring the misfit of the analysis to the background, the data and certain a priori constraints.

$$J = \lambda_{\text{CONV}} J_{\text{CONV}} + \lambda_{\text{SCAT}} J_{\text{SCAT}} + \lambda_{\text{SPD}} J_{\text{SPD}} + \lambda_{\text{VWM}} J_{\text{VWM}} + \lambda_{\text{LAP}} J_{\text{LAP}} + \lambda_{\text{DIV}} J_{\text{DIV}} + \lambda_{\text{VOR}} J_{\text{VOR}} + \lambda_{\text{DYN}} J_{\text{DYN}}$$


Term	Expression	Description of constraint
		Observation Function for the
J_{CONV}	$\sum (\mathbf{V}_A - \mathbf{V}_O)^2$	• wind vectors
J_{SCAT}	$\sum (\mathbf{V}_A - \mathbf{V}_O)^2$	• wind vectors
J_{SPD}	$\sum (\mathbf{V}_A - \mathbf{V}_O)^2$	• wind speeds
		Background Constraints on the
J_{VWM}	$\int (\mathbf{V}_A - \mathbf{V}_B)^2$	• vector wind magnitude
J_{LAP}	$\int [\nabla^2 (u_A - u_B)]^2 + \int [\nabla^2 (v_A - v_B)]^2$	• Laplacian of the wind components
J_{DIV}	$\int [\nabla^2 (\chi_A - \chi_B)]^2$	• divergence
J_{VOR}	$\int [\nabla^2 (\psi_A - \psi_B)]^2$	• vorticity
J_{DYN}	$\int (\partial \zeta_A / \partial t - \partial \zeta_B / \partial t)^2$	• vorticity tendency

$$\mathbf{V}_A = \alpha \mathbf{V}_A + \mathbf{V}_\delta$$


Processing Status



	YEAR																									
	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10	11	12
F08	█	█	█	█	█																					
F10				█	█	█	█	█	█	█	█	█														
F11					█	█	█	█	█	█	█	█	█	█												
F13										█	█	█	█	█	█	█	█	█	█	█	█	█	█	█	█	█
F14											█	█	█	█	█	█	█	█	█	█	█	█	█	█	█	█
TMI											█	█	█	█	█	█	█	█	█	█	█	█	█	█	█	█
F15													█	█	█	█	█	█	█	█	█	█	█	█	█	█
QSCAT													█	█	█	█	█	█	█	█	█	█	█	█	█	█
AMSRE																█	█	█	█	█	█	█	█	█	█	█
Seawinds																	█									




Outputs








- Three products are produced for distribution to the community at 25km resolution:
 - Level3.0: 6-hourly global analyses (U,V)
 - Level3.5: 5-day and monthly means (U,V,W,USTR,VSTR)
 - 3.5a: satellite only
 - 3.5b: all data
 - Level2.5: Passive satellite wind speed observations (SSM/I, AMSRE,TMI) with directions assigned from the analysis (U,V)

***All products reside on the Wentz "bytemap" 25km grid.*





Validation




- Examine analysis fit to assimilated observations (rms, mean speed and direction differences)
- Examine the analysis fit to independent observations (eg Windsat, NSCAT, ERS)
- Compare VAM analyses to operational analyses, satellite imagery, and climatologies.




Statistics :: all data used






ECMWF 10m Winds




06Z02JAN2004

VAM FLK V1.1 Winds




06Z02JAN2004




ECMWF Analysis

VAM FLK V1.0 Analysis



06Z02JAN2004



06Z02JAN2004

NWP Impact experiments



Datasets

QSCAT SLP (QSLP)
QSCAT Gradient Level Winds (QGLW)
QSCAT Sea Level Winds (QSLW)

ASCAT SLP (ASLP)
ASCAT Gradient Level Winds (AGLW)
ASCAT Sea Level Winds (ASLW)

SSM/I, TRMM TMI, AMSRE, Windsat

Global Experiments

GEOS-5 (QSLW)	NCEP (QSLW)
GEOS-5 (QSLP)	
GEOS-5 (QGLW)	VAM (QSLW)
GEOS-5 (All QSCAT)	VAM (ASLW)
	VAM (All SSWs)
GEOS-5 (ASLW)	
GEOS-5 (ASLP)	
GEOS-5 (AGLW)	
GEOS-5 (All ASCAT)	

Regional Experiments

WRF (QSLW)	WRF (ASLW)
WRF (QSLP)	WRF (ASLP)
WRF (QGLW)	WRF (AGLW)
WRF (All QSCAT)	WRF (All ASCAT)
WRF (All Datasets)	



Regional Impact Studies

2005	2008
Katrina (8/23-8/30)	Gustav (8/25-9/4)
Ophelia (9/6-9/18)	Hanna (8/28-9/7)
Phillipe (9/17-9/23)	Ike (9/1-9/15)
2006	
Ernesto (8/24-9/3)	
Gordon (9/11-9/20)	
Helene (9/12-9/24)	

Impact of Seawinds on VAM Analysis



Ambiguous

Unique




NCEP Analysis (interp to .5deg,3hrs)

VAM Analysis




April 11, 2003 3Z


Impact of Seawinds on VAM Vorticity Analysis




**NCEP
Analysis**



**VAM
Analysis**



Difference




April 11, 2003 3Z


Example of a Cyclone Added by Seawinds in the North Atlantic

(shaded field represents clouds from GOES East IR4)


NCEP Analysis




VAM Analysis



Ambiguous Seawinds



Unique Seawinds




April 11 2003 00Z


Example of an Anticyclone Deleted by Seawinds in the South Atlantic

(shaded field represents clouds from GOES East IR4)


NCEP Analysis




VAM Analysis



Ambiguous Seawinds



Unique Seawinds



April 11 2003 00Z




Figure 6 Impact of Quikscat winds on the NCEP and GEOS-5 surface wind analysis at the initial insertion time (top), 6-hours into the assimilation (middle) and 1-week (bottom). Shaded colors indicate the magnitude of the vector difference between the control and control+quikscat experiments.







Figure 10 RMS fit of the NCEP and GEOS-5 analyses to Quikscat for July 2005 and September 2006 respectively. The NCEP and ECMWF operational analyses are shown for comparison. The recently created Cross-Calibrated Multi-Platform (CCMP) surface wind analysis is also shown. The CCMP data set represents the best fit to Quikscat since it is a high resolution analysis (25 km) with less constraints for smoothness.



Analysis	Cyclones Added	Cyclones Deleted	Position Impact		Vorticity Impact		Max Wind Impact	
			Avg (km)	Max	Avg	Max	Avg (m/s)	Max
NCEP	155	346	89	186	-0.4	2.4	0.5	2.4
GEOS-5	309	379	100	251	0.1	4.5	0.7	3.0






Figure 12 NCEP 1000mb wind analysis on September 1, 2005 6Z. The top left panel shows the NCEP Control analysis (no Quikscat) depicting a closed circulation center (located at the endpoint of the vector marker). Quikscat wind barbs (red) are show for reference. The top right panel shows the 1000mb wind analysis for the NCEP analysis that assimilated the Quikscat winds. The NCEP operational analysis (bottom left) is shown for comparison. In general, the assimilation of Quikscat in the NCEP DAS results in many more deleted cyclones than added as can be seen in this example.

Estimation of Observation Impact on NWP




24h Forecast Error (J/Kg)



- control
- no airs
- no raob
- no amsua
- no qkscat
- no satwind

Observation Count (millions)


24h Forecast Error Reduction (J/Kg)



NASA
GEOS-5 July
2005 00z

Gelaro, Zhu, ...

Sea Level Pressure S Hem Extra Tropics




g5_068 - GEOS-5 2.1.4 Half-degree Control
g5_069 - GEOS-5 2.1.4 Half-degree ASCAT
g5_070 - GEOS-5 2.1.4 Half-degree QSCAT

Average of 22 Five-Day Forecasts

g5_068 vs ncep
g5_069 vs ncep
g5_070 vs ncep



1-week Impact on GEOS-5 Analysis of Winds in OSSE





Perfect QSLW

Perfect QSLP



200mb






500mb




850mb




10m





1-week Impact on GEOS-5 Analysis of SLP in OSSE



Perfect QSLW



Perfect QSLP





1-week Impact on GEOS-5 Analysis of Heights in OSSE



Perfect QSLW

Perfect QSLP




200mb





500mb





850mb




GEOS-5 Global Analysis Error (Day-1) in OSSE



GEOS-5 Global Analysis Error (Day-7) in OSSE





GEOS-5 Analysis of SLP and 10m Winds in OSSE





Control

QSLW





QSLP

Nature



September 18, 1999 06Z

GEOS-5 Analysis of Winds in OSSE




September 18, 1999 06Z



NWP Summary

- The assimilation of Quikscat winds improves NASA and NCEP model analyses of ocean sfc wind; the impact on model forecasts is positive but smaller than previously obtained. The impact is limited by competing data, superobing, and various aspects of the GSI analysis.
- The in-depth analysis is ongoing, with a focus on the influence of QuikSCAT data on analyses and forecasts of surface winds, and the vertical dynamic and thermodynamic structure.
- Modifications to quality control, error specification, and data thinning are being tested.
- Quikscat derived sea level pressure and gradient level winds are being assimilated using the GEOS-5, NCEP and WRF models in combination with the Quikscat surface winds to assess the potential for increased beneficial impact of Quikscat as mass and wind data.
- Quikscat data were used successfully in the development of the 1/8 and 1/12 degree versions of the fvGCM, and this application will continue as we go to even higher model resolution. (Atlas et al., 2005, 2007; Shen et al., 2006a,b).

CCMP Summary



- 20+ years of CCMP ocean surface wind velocity data at 6 hour intervals and 25 km resolution have been generated.
- The VAM analyses fit the cross-calibrated data sets very closely with significant improvements in the location and structure of meteorological features.
- The infrastructure is in place to rapidly reprocess based on recommendations from the OVW community with the goal of producing the best possible product.